


Name of Child: \_\_\_\_\_ Date: \_\_\_\_\_

# Clean Intermittent Catheterization (CIC) For Males

## What it is:

**Clean intermittent catheterization (CIC)** is a clean, but non-sterile, way to put a rubber or plastic tube (**catheter**) into the urinary opening (**urethra**) to empty the bladder.



A catheter.

## Why it is done:

Some medical conditions can cause a child to be unable to empty the bladder. When urine remains in the bladder for a long time, bacteria can grow rapidly. CIC helps prevent urinary tract infections by emptying the bladder completely and keeping the over-full bladder from stretching. When CIC is done every three to six hours, the bladder will not over-fill. This may also help prevent dribbling and wetting. CIC can help children to be more active and have a better self-image by helping them stay dry.

## Where it is done:

You can do CIC in any regular toilet facility or private area. Boys may stand, sit or lie down. Consider the child's comfort when you choose a place for CIC.


# The urinary system:

The kidney filters waste products from the whole body. Waste goes into the urine. Urine drains into the bladder, and is stored there until it leaves the body. When urine drains well into the bladder, the kidneys are protected from infection.

## When to catheterize:

To drain the urine well and prevent infection, do CIC regularly.

Your doctor will tell you how often to catheterize your child. As your child gets older, he can learn to do CIC himself. He should follow the same steps you do.



### Write the times your child needs to be catheterized here:

- first thing in the morning
- every \_\_\_\_\_ hours during the day
- when my child goes to bed
- when I go to bed
- other times: \_\_\_\_\_

If your child has his last drink two hours before bedtime, it may help him stay dry through the night.

Do not wake up a baby at night to do CIC. Instead, catheterize when you and your baby are awake for feedings.


## Catheters come in many sizes:

Here are the sizes used by most people, based on age:

<b>Age</b>	<b>Catheter Size</b>
newborns	5 French
infants	8 French
toddlers and young school-aged	10 French
preteens	12 French
adolescents	14 French
older adolescents and adults	16 French

## To prevent infections:

To prevent infections, have your child drink lots of fluids, wash your hands carefully before and after catheterization, keep supplies clean and keep the area around the urinary opening clean.




Clean your hands with soap and water.

## How to do Clean Intermittent Catheterization:

### What you need:

- catheter, size \_\_\_\_\_ French
- something to store the catheter in when not using it, like a toothbrush holder or clean plastic bag
- soap and water, soap towelettes, or cotton balls with soap
- urine container
- water-soluble lubricant (like K-Y Jelly)
- clean gloves (If your child can do CIC without help, you do not need gloves.)



A catheter.

### What to do:

It is best to have your child help with clean intermittent catheterization as much as he is able, while you watch. This gives your child independence, control, and a better self-image. Explain to your child what you are doing as you are doing it.

1. Have all the supplies you need near and ready to use.
2. Clean your hands with soap and water for 15 to 30 seconds. Rinse, then dry your hands with clean paper towels.

If your hands are not visibly dirty, you can use alcohol-based hand rub to clean them using the following steps:

1. Put enough alcohol-based hand rub to fill the palm of your hand.
2. Spread the hand rub over all parts of your hands.
3. Rub hands until dry.

3. Put on clean gloves.

4. Put your child in position for catheterization.


5. Arrange your child's clothes so they do not get wet.



**Clean gloves.**

6. Put water-soluble lubricant on the first two inches of the catheter.

7. Hold your child's penis straight. If your child is not circumcised, pull back the foreskin as far as it will go without forcing it.




**Put water-soluble lubricant on the first two inches of the catheter.**

8. Use the towelettes or soapy cotton balls to wash around the urinary opening (urethra). Begin at the opening and wash outward in a circular motion. Repeat this two more times, using a clean towelette or cotton ball each time.

9. Put the urine container in place to catch the urine, or have the child near the toilet.

10. Gently slide the catheter into the urinary opening until urine starts to flow. Never force the tube. If you have trouble passing the tube, stop and call your child's doctor.




11. Let all the urine flow out either into the urine container or the toilet.

12. Have the child bear down a few times with the catheter still in place. This helps to empty all the urine from the bladder.

13. Slowly start to take out the catheter. If urine starts to flow again, stop pulling out the catheter until the flow of urine stops.

14. Pinch the catheter, and keep it pinched while you take it out. This keeps urine from flowing back into the bladder.
15. After you take out the catheter, wipe across the urinary opening with a towelette or toilet paper. This will prevent odors. If your child is not circumcised, pull the foreskin back over the head of the penis when you are finished.
16. Praise your child for his help, and return his clothing to its normal position.
17. Wash the catheter with soap and water. Rinse the inside of the catheter with clear water, dry it, and put it back in the clean, dry container.
18. If you used a urine container, empty it in a toilet. Wash and dry the container.
19. Take off the gloves, and throw them out. Wash and dry your hands.
20. If your doctor wants you to keep records, write down the time and amount of urine.



**Write down the time and amount of urine.**

## **Problem-Solving:**

- If urine leaks around the catheter, try using a larger-sized catheter.
- If the tip of the catheter will not slide into the urinary opening, do not force it. Try using a smaller-sized catheter.
- Bleeding from the urinary opening can be caused by injury during the catheterization. If this occurs, stop the catheterization and call your doctor.
- Trouble passing the catheter can be caused by a tight muscle (increased sphincter tone) due to anxiety or spasm. Change the position of the penis and push gently on the catheter until the muscle relaxes. If you still cannot pass the catheter, call your doctor.
- No urine on catheterization may mean the catheter is not in the right place. Check the placement of the catheter.

## Call your child's doctor if you see signs of a urinary tract infection:

- blood in urine
- chills
- increased wetting
- fever
- bladder spasms
- pain or tenderness over lower back



**Call your doctor  
if you see signs of  
a urinary tract infection.**

## Care of the Catheter:

### What you need:

- catheter
- something to keep the catheter in, like a toothbrush holder or clean plastic bag
- soap
- clean gloves
- tap water
- a clean container for water

### What to do:

- Wash the catheter with soap and water. Rinse the inside of the catheter with clear water. Let it dry before you store it in a clean container.
- Every night, let the catheter air dry on a clean surface.
- Every night, wash the container you use to store and carry the catheter. Use soap and water, and let it air dry. If you use a plastic sandwich bag to store the catheter, use a new bag each day.
- One catheter can last one to three months. If the catheter becomes too soiled to use, or becomes difficult to put in, throw it out.

Take the catheter and supplies everywhere you go. It is important to stick to the catheterization schedule to avoid infections and wetting. CIC can help children be normally active and have a better self-image by helping them stay dry.

## Now that you've read this:

- Show your nurse or doctor how you will catheterize your child. (Check when done.)
- Show your nurse or doctor how you will keep the catheter clean. (Check when done.)
- Tell your nurse or doctor when you need to call the doctor's office. (Check when done.)
- yes  no See the handout: **Latex Allergy #249**



If you have any questions or concerns,  
 call your child's doctor or  call \_\_\_\_\_

If you want to know more about child health and illness,  
visit our library at The Emily Center at Phoenix Children's Hospital  
1919 East Thomas Road  
Phoenix, AZ 85016  
602-933-1400  
866-933-6459  
[www.phoenixchildrens.org](http://www.phoenixchildrens.org)  
Facebook: [facebook.com/theemilycenter](https://facebook.com/theemilycenter)  
Twitter: [@emilycenter](https://twitter.com/emilycenter)  
Pinterest: [pinterest.com/emilycenter](https://pinterest.com/emilycenter)

**Disclaimer**

The information provided at this site is intended to be general information, and is provided for educational purposes only. It is not intended to take the place of examination, treatment, or consultation with a physician. Phoenix Children's Hospital urges you to contact your physician with any questions you may have about a medical condition.

November 17, 2016 • DRAFT to family review  
#1731 • Written by Barbara Singer, MS, RN • Updated by Gwen Grimbsy, MD  
Illustrated by Dennis Swain

## Clean Intermittent Catheterization (CIC) For Boys

Name of Health Care Provider: \_\_\_\_\_

Date returned: \_\_\_\_\_  db

### Family Review of Handout

Health care providers: Please teach families with this handout.

Families: Please let us know what you think of this handout.

Would you say this handout is hard to read?  Yes  No

easy to read?  Yes  No

Please circle the parts of the handout that were hard to understand.

Would you say this handout is interesting to read?  Yes  No

Why or why not?

Would you do anything differently after reading  
this handout?  Yes  No

If yes, what?

After reading this handout, do you have any  
questions about the subject?  Yes  No

If yes, what?

Is there anything you don't like about the drawings?

Yes

No

If yes, what?

What changes would you make in this handout to make it better or easier to understand?

Please return your review of this handout to your nurse or doctor or send it to the address below.

The Emily Center  
Health Education Specialist  
Phoenix Children's Hospital  
1919 East Thomas Road  
Phoenix, AZ 85016-7710

602-933-1395

**Thank you for helping us!**